

A LOWER BOUND FOR PARTIAL LIST COLORINGS

GLENN G. CHAPPELL

Department of Mathematics, Southeast Missouri State University

ABSTRACT. Let G be an n -vertex graph with list-chromatic number χ_ℓ . Suppose each vertex of G is assigned a list of t colors. Albertson, Grossman, and Haas [1] conjecture that at least $\frac{tn}{\chi_\ell}$ vertices can be colored from these lists. We prove a lower bound for the number of colorable vertices. As a corollary, we show that at least $\frac{6}{7}$ of the conjectured number can be colored.

1. INTRODUCTION

Let G be an n -vertex graph. Let each vertex of G be assigned a list of s colors. An s -list coloring of G is a proper vertex coloring in which each vertex is given a color from its list. We say G is s -choosable if G has an s -list coloring for each assignment of lists of size s to the vertices. The *list-chromatic number* of G , denoted by χ_ℓ , is the least s so that G is s -choosable. List coloring was introduced independently by Vizing [4] and by Erdős, Rubin, and Taylor [2].

Suppose each vertex of G is assigned a list of t colors ($t \leq \chi_\ell$). It may not be possible to color every vertex from these lists; however, we can color a subset of the vertices. Albertson, Grossman, and Haas [1] ask how many vertices can be colored. Letting ℓ_t range over all assignments of lists of size t to the vertices of G , we define

$$\lambda_t := \min_{\ell_t} \{ \text{max. no. of vertices of } G \text{ colorable from the lists } \ell_t \}.$$

Conjecture 1.1 (Albertson, Grossman, & Haas [1]). *If $t \leq \chi_\ell$, then $\lambda_t \geq \frac{tn}{\chi_\ell}$. $\square$*

Conjecture 1.1 clearly holds when $t = 0$ or $t = \chi_\ell$. When $t = 1$, we note that G has an independent set of size at least $\frac{n}{\chi} \geq \frac{n}{\chi_\ell}$, where χ is the ordinary chromatic number. Since an independent set can always be colored from lists of size 1, we have $\lambda_1 \geq \frac{n}{\chi_\ell}$, and so the conjecture holds for $t = 1$. However, the conjecture remains open for $1 < t < \chi_\ell$.

Date: May 13, 1998.

1991 Mathematics Subject Classification. 05C15.

We will use a method similar to that of the proof of [1, Thm. 2] to prove a lower bound for λ_t . As a corollary, we show that $\lambda_t > \frac{6}{7} \cdot \frac{tn}{\chi_\ell}$.

2. THE RESULTS

For integers $s > t > 0$, we define

$$f_{s,t}(x) := 1 - x - \left[1 - \frac{1-x}{s-t}\right]^t.$$

Then $f_{s,t}(0) > 0$, $f_{s,t}(1) < 0$, and $\frac{d}{dx}f_{s,t}(x) < 0$ for $x > 0$. Thus, the equation $f_{s,t}(x) = 0$ has exactly one positive solution, which we denote by $q_{s,t}$. Note that $0 < q_{s,t} < 1$.

Theorem 2.1. *If G is s -choosable, and $s > t > 0$, then $\lambda_t \geq q_{s,t} \cdot n$. Furthermore, this inequality is strict if $t > 1$.*

Proof. Let $q = q_{s,t}$. Let each vertex of G be assigned a list of t colors, and let R be the union of these lists. Let $u = s - t$. We augment each list by adding new colors $\pi_1, \pi_2, \dots, \pi_u \notin R$. There is an s -list coloring φ from the augmented lists. For $1 \leq i \leq u$, let I_i be the set of vertices given color π_i ; each I_i is an independent set. Let $\mathcal{I} = I_1 \cup \dots \cup I_u$, and let $H = G - \mathcal{I}$. Thus, φ colors the vertices of H with colors in R .

We partition the set R randomly into classes $R_0, R_1, \dots, R_u$; the class in which each color falls is chosen independently. We place a given color in class R_0 with probability q and in each of the other classes with probability $\frac{1-q}{u}$.

Given a partition of R , we construct a proper t -list coloring of a subset of the vertex set of G as follows. The colors in R_0 are used on the vertices of H . We color a vertex in H with its color in the coloring φ if that color appears in R_0 ; otherwise, we leave the vertex uncolored. Thus, each vertex in H is colored with probability q .

For $1 \leq i \leq u$, the colors in R_i are used on I_i . We color a vertex in I_i if a color in its list appears in R_i ; otherwise, we leave the vertex uncolored. A particular color fails to appear in R_i with probability

$$1 - \frac{1-q}{u}.$$

Thus, each vertex of $\mathcal{I}$ is colored with probability

$$1 - \left[1 - \frac{1-q}{u}\right]^t = f_{s,t}(q) + q = q,$$

since $f_{s,t}(q) = 0$.

For some partition of R the number of colored vertices is at least the expected number. Since the above scheme results in each vertex being colored with probability q , we have $\lambda_t \geq qn$.

Now let $t > 1$. To see that the above inequality is strict, note that q is a root of the polynomial $p(x) := u^t f_{s,t}(x)$. All the coefficients of $p(x)$ are integers; the leading coefficient of is -1 . By the Rational Roots Theorem, every rational root of $p(x)$ is an integer. Since $0 < q < 1$, we conclude that q is irrational, and so $\lambda_t \neq qn$. $\square$

Albertson et al. showed [1, Thm. 2] that $\lambda_2 > \frac{3n}{5}$ if $\chi_\ell = 3$. Theorem 2.1 allows us to improve on this. Since $q_{3,2} = \frac{-1+\sqrt{5}}{2}$, we have the following stronger result.

Corollary 2.2. *If $\chi_\ell = 3$, then $\lambda_2 > \frac{-1+\sqrt{5}}{2} \cdot n$. $\square$*

The conjectured lower bound is greater still: $\frac{2n}{3}$.

Remark 2.3. Thomassen [3] showed that every planar graph is 5-choosable. Thus, we can apply Theorem 2.1 to planar graphs. Albertson et al. [1] estimate that their method could be used to show that $\lambda_4 \geq \frac{7}{10}n$ for every planar graph. By definition, $q_{5,4}$ is the positive root of $1 - x - x^4$: a bit more than 0.724. Thus, $\lambda_4 > 0.724n$ if $\chi_\ell \leq 5$ and in particular for every planar graph. $\square$

Next we show that λ_t is at least $\frac{6}{7}$ of the conjectured lower bound.

Corollary 2.4. *If $t \leq \chi_\ell$, then $\lambda_t > \frac{6}{7} \cdot \frac{tn}{\chi_\ell}$. $\square$*

Corollary 2.4 follows from Theorem 2.1, the comments after Conjecture 1.1, and the following lemma.

Lemma 2.5. *If $s > t > 0$, then $\frac{6}{7} \cdot \frac{t}{s} < q_{s,t} \leq \frac{t}{s}$.*

Proof. We first show that

$$\frac{6}{7} \cdot \frac{t}{s} < q_{s,t}.$$

Recall that $f_{s,t}(q_{s,t}) = 0$, and that $f_{s,t}(x)$ is decreasing for $x > 0$. Thus, it suffices to show that $f_{s,t}(cv) > 0$, where $c = \frac{6}{7}$ and $v = \frac{t}{s}$.

$$\begin{aligned} f_{s,t}(cv) &= 1 - cv - \left[1 - \frac{1 - cv}{s - sv} \right]^{sv} \\ &> 1 - cv - e^{-v \cdot \frac{1 - cv}{1 - v}}, \quad \text{since } \left(1 - \frac{a}{b} \right)^b < e^{-a}, \text{ for } 0 < a < b. \end{aligned}$$

We show that the last expression is positive. Note that both $1 - cv$ and $e^{-v \cdot \frac{1 - cv}{1 - v}}$ are positive; let $g(v)$ be the difference of the logarithms of these quantities.

$$g(v) := \ln(1 - cv) - \left(-v \cdot \frac{1 - cv}{1 - v} \right).$$

It suffices to show that $g(v) > 0$, for $0 < v < 1$. Verification of this requires only the standard techniques of calculus and is omitted.

Now we show that

$$q_{s,t} \leq \frac{t}{s}.$$

Since $f_{s,t}(x)$ is decreasing for $x > 0$, it suffices to show that $f_{s,t}\left(\frac{t}{s}\right) \leq 0$.

$$\begin{aligned} f_{s,t}\left(\frac{t}{s}\right) &= 1 - \frac{t}{s} - \left[1 - \frac{1 - \frac{t}{s}}{s - t}\right]^t \\ &= 1 - \frac{1}{s} \cdot t - \left[1 - \frac{1}{s}\right]^t \\ &\leq 0. \quad \square \end{aligned}$$

The techniques of the above proof can be used to prove a slightly better lower bound for $q_{s,t}$. Numerical computations indicate that

$$\inf_{s,t} \frac{q_{s,t}}{t/s} \approx 0.8598841287 > \frac{6}{7}.$$

REFERENCES

- [1] M. O. Albertson, S. Grossman, and R. Haas, Partial List Colorings, manuscript, April 15, 1998.
- [2] P. Erdős, H. Rubin, and A. Taylor, Choosability in Graphs, *Congr. Numer.* **26** (1979), 126–157.
- [3] C. Thomassen, Every Planar Graph is 5-Choosable, *J. Combin. Theory Ser. B* **62** (1994), 180–181.
- [4] V. G. Vizing, Coloring the vertices of a graph in prescribed colors (Russian), *Diskret. Analiz.* **29** (1976), 3–10.

DEPARTMENT OF MATHEMATICS, SOUTHEAST MISSOURI STATE UNIVERSITY,
CAPE GIRARDEAU, MO 63701, USA

E-mail address: gchappell@semovm.semo.edu